

Letters of Intent: EPR
Target Date: February 15, 2003

Oceanographic and topographic influences on dispersal of hydrothermal vent species

L.S. Mullineaux, A.M. Thurnherr (with K.G. Speer as collaborator), J. R. Ledwell, D. McGillicuddy, and J.W. Lavelle

The population dynamics, community structure and gene flow of species living at deep-sea hydrothermal vents depend strongly on migration of their larvae between isolated vent sites. We are interested in quantifying the extent and direction of larval dispersal from vents near 9°N on the East Pacific Rise. We also want to solve the reverse question – i.e., to what extent are EPR vent populations recolonized by their own larvae. Because retention and dispersal of larvae are controlled largely by the oceanic flow field, we propose to investigate these processes using a tracer-release experiment, current meter measurements and a hydrodynamic/transport model. Since some larvae likely modify their vertical positions over the course of their larval stages, which can modify their dispersal in sheared flow, we plan to incorporate vertical larval swimming into our dispersal model. We will infer vertical swimming behaviors by sampling larvae near vents at heights between 1 and 200 mab to determine what layers of the deep water column they inhabit at different stages in their larval lives.

We anticipate that this study will require participation in three cruises over a period of roughly 1 year. The tracers (dye and SF6) will be released on Cruise 1, and the dye surveyed immediately. SF6 will be surveyed at 35 d (Cruise 2) and will be monitored from a mooring for a year at a site to the north of the Clipperton FZ. The current meter array will be deployed on Cruise 1 and recovered on Cruise 3. Vertical distributions of larvae in the plankton will be sampled on each cruise. Modeling efforts will be initiated prior to the cruises and continue after them.

We are hoping to coordinate these oceanographic measurements with studies of chemistry, heat flux or other processes that are influenced by circulation and dispersion at mid-ocean ridges.